

02/881754


PCT/EP

99/08744

Europäisches
PatentamtEuropean
Patent OfficeOffice européen
des brevets

ESU

EP 93/8744



Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98121478.6

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN
THE HAGUE, 11/01/00
LA HAYE, LE

THIS PAGE BLANK (USPTO)



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.: 98121478.6
Demande n°:

Anmeldetag:
Date of filing: 12/11/98
Date de dépôt:

Anmelder:
Applicant(s):
Demandeur(s):
Nitsch, Roger M., Prof. Dr.
20146 Hamburg
GERMANY

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:
Methods of diagnosing or treating neurological diseases

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C12N15/12, C07K14/47, C12Q1/68, G01N33/68, A61K38/17

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

THIS PAGE BLANK (USPTO)

12-11-1998

EP98121478.6

SPEC

METHODS OF DIAGNOSING OR TREATING NEUROLOGICAL DISEASES

Neurological diseases are widely spread within a population and have a strong impact not only on patients' life but also on society as such. Therefore, there is a strong need to elucidate the causes and the underlying pathogenesis of such neurological diseases. Among such neurological diseases, Alzheimer's disease (AD) has a predominant position. Alzheimer's disease, first described by the Bavarian psychiatrist Alois Alzheimer in 1907, is a progressive neurological disorder which begins with short term memory loss and proceeds to loss of cognitive functions, disorientation, impairment of judgment and reasoning and, ultimately, dementia. It is the most common cause of dementia. The neuropathology is characterized by the formation in brain of amyloid plaques and neurofibrillary tangles. AD has been estimated to afflict 5 to 11 percent of the population over age 65 and as much as 47 percent of the population over age 85. Moreover, as adults, born during the population boom of the 1940's and 1950's, approach the age when AD becomes more prevalent, the control and treatment of AD will become an even more significant health care problem. Familial forms of AD are genetically heterogeneous, but most with early onset are linked to mutations in the presenilin genes *PSEN1* and *PSEN2*, as well as to mutations of the amyloid precursor gene *APP*. The majority of AD patients have no obvious family history and are classified as sporadic AD.

As neurological diseases, in particular AD, are a growing social and medical problem, there is a strong need for methods of diagnosing or prognosing said diseases in subjects as well as for methods of treatment.

In one aspect, the invention features a protein molecule (hereinafter "SELADIN-1") shown in SEQ ID NO. 1.

In another aspect, the invention features an isolated nucleic acid molecule encoding SELADIN-1 shown in SEQ ID NO. 1. Said nucleic acid molecule can be a DNA molecule, such as a genomic DNA molecule or a cDNA molecule, or an RNA molecule, such as a mRNA molecule. In particular, said nucleic acid molecule can be a cDNA

12-11-1998

EP98121478.6

SPEC

2

molecule having a coding sequence substantially the same as the coding sequence of SEQ ID NO. 2.

In another aspect, the invention features an oligonucleotide which hybridizes with a base sequence encoding SELADIN-1 shown in SEQ ID NO.1 in a solution of 0.2xSSC (standard saline citrate) and 0.1 % SDS at 60 °C.

In another aspect, the invention features an isolated DNA comprising a base sequence which hybridizes with said base sequence encoding SELADIN-1 shown in SEQ ID NO. 1 in a solution of 50 % formamide, 4xSSC, 50 mM HEPES, pH 7.0, 10x Denhardt's solution, 100 µg/ml thermally denatured salmon sperm DNA at 42 °C. The invention further aims at an oligonucleotide which hybridizes with a base sequence comprising the afore mentioned DNA molecule in a solution of 0.2xSSC and 0.1 % SDS at 60 °C.

"Variants" of a protein molecule shown in SEQ ID NO.1 include e.g. proteins with conservative amino acid substitutions in highly conservative regions. For example, isoleucine, valine and leucine can each be substituted for one another. Aspartate and glutamate can be substituted for each other. Glutamine and asparagine can be substituted for each other. Serine and threonine can be substituted for each other. Amino acid substitutions in less conservative regions include e.g.: Isoleucine, valine and leucine can each be substituted for one another. Aspartate and glutamate can be substituted for each other. Glutamine and asparagine can be substituted for each other. Serine and threonine can be substituted for each other. Glycine and alanine can be substituted for each other. Alanine and valine can be substituted for each other. Methionine can be substituted for each of leucine, isoleucine or valine, and vice versa. Lysine and arginine can be substituted for each other. One of aspartate and glutamate can be substituted for one of arginine or lysine, and vice versa. Histidine can be substituted for arginine or lysine, and vice versa. Glutamine and glutamate can be substituted for each other. Asparagine and aspartate can be substituted for each other.

"Variants" of SELADIN-1 shown in SEQ ID NO.1 also include .g. protein molecules which are encoded by DNA fragments comprising base sequences which hybridiz with

said base sequence encoding SELADIN-1 shown in SEQ ID NO. 1 in a solution of 50 % formamide, 4xSSC, 50 mM HEPES, pH 7.0, 10x Denhardt's solution, 100 µg/ml thermally denatured salmon sperm DNA at 42 °C.

The invention further features an antibody specifically immunoreactive with an immunogen, wherein said immunogen is SELADIN-1 shown in SEQ ID NO. 1 or a variant or fragment thereof. In another aspect, the invention aims at a method of detecting pathological structures in the brain of a subject which comprises immunocytochemically staining said pathological structures with said antibody.

In another aspect, the invention features a vector comprising the isolated nucleic acid encoding a SELADIN-1 shown in SEQ ID NO. 1 or a variant thereof. In preferred embodiments, a virus, a bacteriophage, or a plasmid comprises said nucleic acid.

In still another aspect, the invention features a plasmid adapted for expression in a bacterial cell which comprises a nucleic acid molecule encoding SELADIN-1 shown in SEQ ID NO. 1 or a variant thereof, and the regulatory elements necessary for expression of said molecule in the bacterial cell.

In a further aspect, the invention features a plasmid adapted for expression in a yeast cell which comprises a nucleic acid molecule encoding SELADIN-1 shown in SEQ ID NO. 1 or a variant thereof, and the regulatory elements necessary for expression of said molecule in the yeast cell.

In another aspect, the invention features a plasmid adapted for expression in a mammalian cell which comprises a nucleic acid molecule encoding SELADIN-1 shown in SEQ ID NO.1 or a variant thereof, and the regulatory elements necessary for expression of said molecule in the mammalian cell.

In a further aspect, the invention features a cell transformed with a nucleic acid molecule encoding SELADIN-1 shown in SEQ ID NO. 1 or a variant thereof. In preferred

embodiments, said cell is a bacterial cell, a yeast cell, a mammalian cell, or an insect cell.

In a further aspect, the invention features a bacterial cell comprising a plasmid adapted for expression in a bacterial cell, said plasmid comprises a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, and the regulatory elements necessary for expression of said molecule in the bacterial cell.

In a further aspect, the invention features a yeast cell comprising a plasmid adapted for expression in a yeast cell, said plasmid comprises a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, and the regulatory elements necessary for expression of said molecule in the yeast cell.

In another aspect, the invention features a mammalian cell comprising a plasmid adapted for expression in a mammalian cell, said plasmid comprises a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, and the regulatory elements necessary for expression of said molecule in the mammalian cell.

In preferred embodiments, said subjects suffer from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

In another aspect, the invention features a method of diagnosing or prognosing a neurological disease in a subject comprising:

determining a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO. 1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, a variant or fragment thereof,

d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),

) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c),

in a sample from said subject,

and

comparing said level, or said activity, or both said level and said activity, of at least one of said substances a) to e) to a reference value representing a known disease or health status,

thereby diagnosing or prognosing said neurological disease in said subject.

In preferred embodiments, said subjects suffer from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

It is particularly preferred that said sample is a brain tissue or other body cells including blood cells and skin fibroblasts. According to the present invention, a reduction in the level, or activity, or both said level and said activity, of SELADIN-1 or SELADIN-1 transcripts in said subject's brain regions affected heavily by neurodegeneration relative to a reference value indicates a diagnosis or prognosis of Alzheimer's disease. Predominantly neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus and the amygdala degenerate in Alzheimer's disease.

It is also particularly preferred that said sample is cerebrospinal fluid or another body fluid.

It might be preferred that said subject has previously been determined to have one or more factors indicating that such subject is afflicted with a neurological disease.

In preferred embodiments, said subject can be a human, an experimental animal, e.g. a rat or a mouse, a domestic animal, or a non-human primate, e.g. a monkey. The experimental animal can be an animal model for a disorder, e.g. a transgenic mouse with an Alzheimer's-type neuropathology.

In preferred embodiments, at least one of said substances is detected using an immunoassay, an enzyme activity assay and/or a binding assay.

In preferred embodiments, measurement of the level of transcription products of the *SELADIN-1* gene, or a variant thereof, is performed in body cells using Northern blots with probes specific for the *SELADIN-1* gene or said variant. Quantitative PCR with primer combinations to amplify *SELADIN-1* gene-specific sequences from cDNA obtained by reverse transcription of RNA extracted from body cells of a subject can also be applied. These techniques are known to those of ordinary skill in the art (see e.g. Watson et al., *Rekombinierte DNA*, 2nd edition, Spektrum Akademischer Verlag GmbH, Heidelberg, 1993; Watson et al., *Recombinant DNA*, 2nd ed. W.H. Freeman and Company, 1992).

In preferred embodiments, said level or activity of the protein molecule shown in SEQ ID NO. 1, or a variant or fragment thereof, is detected using an immunoassay. These assays can measure the amount of binding between said protein molecule ("SELADIN-1") and an anti-SELADIN-1 antibody by the use of enzymatic, chromodynamic, radioactive, or luminescent labels which are attached to either the anti-SELADIN-1 antibody or a secondary antibody which binds the anti-SELADIN-1 antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

The antibody or ligand to be used should preferably specifically detect SELADIN-1 or a variant or fragment thereof. It is preferred that it does not substantially interact with any other protein present in said sample.

Monoclonal antibodies capable of recognizing a protein molecule of SEQ ID NO. 1 or a variant or fragment thereof can be prepared using methods known in the art (see e.g. Köhler and Milstein, Nature 256, 495 - 497 1975; Kozbor et al., Immunol. Today 4, 72, 1983; Cole et al., Monoclonal antibodies and cancer therapy, Alan R. Liss, Inc., pp 77 - 96, 1985; Marks et al., J. Biol. Chem., 16007 - 16010, 1992; the contents of which are incorporated herein by reference). Such monoclonal antibodies or fragments thereof can also be produced by alternative methods known to those of skill in the art of recombinant DNA technology (see e.g. Sastry et al, PNAS 86: 5728, 1989; ; Watson et al., Rekombinierte DNA, 2nd ed., Spektrum Akademischer Verlag GmbH, 1993; Watson et al, Recombinant DNA, 2nd ed., W. H. Freeman and Company, 1992; the contents of which are incorporated herein by reference). Monoclonal antibodies useful in the methods of the invention are directed to an epitope of SELADIN-1 or a variant or fragment thereof, such that the complex formed between the antibody and SELADIN-1, or between the antibody and said variant or fragment, can be recognized in detection assays. The term "antibodies" encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, single chain antibodies as well as fragments thereof which specifically bind to SELADIN-1, or to a variant or fragment thereof.

Antibodies or ligands might also be used in detecting specifically molecules mentioned under d) and e) above.

If luminescent labels are used in any detection assay, it is preferred to use a confocal optical set-up.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e) in a sample from a subject not suffering of said neurological disease. The healthy subject can be of the same weight, age, and gender as the subject who is being diagnosed or prognosed for said neurological disease. In some cases, it might be preferred to use a reference value from the subject which is diagnosed.

In another aspect, the invention features a method of monitoring the progression of a neurological disease in a subject, comprising:

determining a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO. 1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
- e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

in a sample from said subject;

and

comparing said level, or said activity, or both said level and said activity, of at least one of said substances a) to e) to a reference value representing a known disease or health status,

thereby monitoring the progression of said neurological disease in said subject.

In a preferred embodiment, the level, or the activity, or both said level and said activity, of at least one of said substances a) to e) in a sample is determined at least twice, e.g. at two points which are weeks or months apart. The levels or activities at these two time points are compared in order to monitor the progression of said neurological disease. It might be preferred to take a series of samples over a period of time. In further preferred embodiments, said subject receives a treatment prior to one or more of said sample gatherings.

In preferred embodiments, said subjects suffer from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration

or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

It is particularly preferred that said sample is a brain tissue or other body cells including blood cells and skin fibroblasts. According to the present invention, a reduction in the level, or activity, or both said level and said activity, of SELADIN-1 or SELADIN-1 transcripts in said subject's brain regions affected heavily by neurodegeneration relative to a reference value indicates a diagnosis or prognosis of Alzheimer's disease. Predominantly neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus and the amygdala degenerate in Alzheimer's disease.

It is also particularly preferred that said sample is cerebrospinal fluid or another body fluid.

It might be preferred that said subject has previously been determined to have one or more factors indicating that such subject is afflicted with a neurological disease.

In preferred embodiments, said subject can be a human, an experimental animal, e.g. a rat or a mouse, a domestic animal, or a non-human primate, e.g. a monkey. The experimental animal can be an animal model for a disorder, e.g. a transgenic mouse with an Alzheimer's-type neuropathology.

In preferred embodiments, at least one of said substances is detected using an immunoassay, an enzyme activity assay and/or a binding assay.

In preferred embodiments, measurement of the level of transcription products of the SELADIN-1 gene, or a variant thereof, is performed in body cells using Northern blots with probes specific for the SELADIN-1 gene or said variant. Quantitative PCR with primer combinations to amplify SELADIN-1 gene-specific sequences from cDNA obtained by reverse transcription of RNA extract d from body cells of a subject can also be applied. These techniques are known to those of ordinary skill in the art (see e.g. Watson et al., Rekombiniertes DNA, 2nd edition, Spektrum Akademischer Verlag GmbH,

Heidelberg, 1993; Watson et al., Recombinant DNA, 2nd ed. W.H. Freeman and Company, 1992).

In preferred embodiments, said level or activity of the protein molecule shown in SEQ ID NO. 1, or a variant or fragment thereof, is detected using an immunoassay. These assays can measure the amount of binding between said protein molecule ("SELADIN-1") and an anti-SELADIN-1 antibody by the use of enzymatic, chromodynamic, radioactive, or luminescent labels which are attached to either the anti-SELADIN-1 antibody or a secondary antibody which binds the anti-SELADIN-1 antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow et al., Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

The antibody or ligand to be used should preferably specifically detect SELADIN-1 or a variant or fragment thereof. It is preferred that it does not substantially interact with any other protein present in said sample.

Monoclonal antibodies capable of recognizing a protein molecule of SEQ ID NO. 1 or a variant or fragment thereof can be prepared using methods known in the art (see e.g. Köhler and Milstein, Nature 256, 495 - 497 1975; Kozbor et al., Immunol. Today 4, 72, 1983; Cole et al., Monoclonal antibodies and cancer therapy, Alan R. Liss, Inc., pp 77 - 96, 1985; Marks et al., J. Biol. Chem., 16007 - 16010, 1992; the contents of which are incorporated herein by reference). Such monoclonal antibodies or fragments thereof can also be produced by alternative methods known to those of skill in the art of recombinant DNA technology (see e.g. Sastry et al, PNAS 86: 5728, 1989; ; Watson et al., Rekombinierte DNA, 2nd ed., Spektrum Akademischer Verlag GmbH, 1993; Watson et al, Recombinant DNA, 2nd ed., W. H. Freeman and Company, 1992; the contents of which are incorporated herein by reference). Monoclonal antibodies useful in the methods of the invention are directed to an epitope of SELADIN-1 or a variant or fragment thereof, such that the complex formed between the antibody and SELADIN-1, or between the antibody and said variant or fragment, can be recognized in detection

assays. The term "antibodies" encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, single chain antibodies as well as fragments thereof which specifically bind to SELADIN-1, or to a variant or fragment thereof.

Antibodies or ligands might also be used in detecting specifically any molecules mentioned under d) to e) above.

If luminescent labels are used in any detection assay, it is preferred to use a confocal optical set-up.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e) in a sample from a subject not suffering of said neurological disease. The healthy subject can be of the same weight, age, and gender as the subject who is being diagnosed or prognosed for said neurological disease. In some cases, it might be preferred to use a reference value from the subject which is diagnosed.

In still another aspect, the invention features a method of evaluating a treatment for a neurological disease, comprising:

determining a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO. 1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, a variant or fragment thereof,
 - d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
 - e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c),
- in a sample from said subject;

12-11-1998

12

and

comparing said lev l, or said activity, or both said level and said activity, of at least one of said substances a) to e) to a reference value representing a known disease or health status,

thereby evaluating said treatment for said neurological disease.

In preferred embodiments, said subjects suffer from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

It is particularly preferred that said sample is a brain tissue or other body cells including blood cells and skin fibroblasts. According to the present invention, a reduction in the level, or activity, or both said level and said activity, of SELADIN-1 or SELADIN-1 transcripts in said subject's brain regions affected heavily by neurodegeneration relative to a reference value indicates a diagnosis or prognosis of Alzheimer's disease. Predominantly neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus and the amygdala degenerate in Alzheimer's disease.

It is also particularly preferred that said sample is cerebrospinal fluid or another body fluid.

It might be preferred that said subject has previously been determined to have one or more factors indicating that such subject is afflicted with a neurological disease.

In preferred embodiments, said subject can be a human, an experimental animal, e.g. a rat or a mouse, a domestic animal, or a non-human primate, e.g. a monkey. The experimental animal can be an animal model for a disorder, e.g. a transgenic mouse with an Alzheimer's-type neuropathology.

In preferred embodiments, at least one of said substances is detected using an immunoassay, an enzyme activity assay and/or a binding assay.

In preferred embodiments, measurement of the level of transcription products of the *SELADIN-1* gene, or a variant thereof, is performed in body cells using Northern blots with probes specific for the *SELADIN-1* gene or said variant. Quantitative PCR with primer combinations to amplify *SELADIN-1* gene-specific sequences from cDNA obtained by reverse transcription of RNA extracted from body cells of a subject can also be applied. These techniques are known to those of ordinary skill in the art (see e.g. Watson et al., *Rekombinierte DNA*, 2nd edition, Spektrum Akademischer Verlag GmbH, Heidelberg, 1993; Watson et al., *Recombinant DNA*, 2nd ed. W.H. Freeman and Company, 1992).

In preferred embodiments, said level or activity of the protein molecule shown in SEQ ID NO. 1, or a variant or fragment thereof, is detected using an immunoassay. These assays can measure the amount of binding between said protein molecule ("SELADIN-1") and an anti-SELADIN-1 antibody by the use of enzymatic, chromodynamic, radioactive, or luminescent labels which are attached to either the anti-SELADIN-1 antibody or a secondary antibody which binds the anti-SELADIN-1 antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

The antibody or ligand to be used should preferably specifically detect SELADIN-1 or a variant or fragment thereof. It is preferred that it does not substantially interact with any other protein present in said sample.

Monoclonal antibodies capable of recognizing a protein molecule of SEQ ID NO. 1 or a variant or fragment thereof can be prepared using methods known in the art (see e.g. Köhler and Milstein, *Nature* 258, 495 - 497 1975; Kozbor et al., *Immunol. Today* 4, 72, 1983; Cole et al., *Monoclonal antibodies and cancer therapy*, Alan R. Liss, Inc., pp 77 - 96, 1985; Marks et al., *J. Biol. Chem.*, 16007 - 16010, 1992; the contents of which are incorporated herein by reference). Such monoclonal antibodies or fragments thereof can

also be produced by alternative methods known to those of skill in the art of recombinant DNA technology (see e.g. Sastry et al, PNAS 86: 5728, 1989; ; Watson et al., Rekombinierte DNA, 2nd ed., Spektrum Akademischer Verlag GmbH, 1993; Watson et al, Recombinant DNA, 2nd ed., W. H. Freeman and Company, 1992; the contents of which are incorporated herein by reference). Monoclonal antibodies useful in the methods of the invention are directed to an epitope of SELADIN-1 or a variant or fragment thereof, such that the complex formed between the antibody and SELADIN-1, or between the antibody and said variant or fragment, can be recognized in detection assays. The term "antibodies" encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, single chain antibodies as well as fragments thereof which specifically bind to SELADIN-1, or to a variant or fragment thereof.

Antibodies or ligands might also be used in detecting specifically any molecules mentioned under d) to e) above.

If luminescent labels are used in any detection assay, it is preferred to use a confocal optical set-up.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e) in a sample from a subject not suffering of said neurological disease. The healthy subject can be of the same weight, age, and gender as the subject who is being diagnosed or prognosed for said neurological disease. In some cases, it might be preferred to use a reference value from the subject which is diagnosed.

In a further aspect, the invention features a kit for diagnosis, or prognosis, of a neurological disease, said kit comprising:

(1) at least one reagent which is selected from the group consisting of reagents that selectively detect

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,

c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,

d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),

e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c), and

(2) instructions for diagnosing, or prognosing said neurological disease by

(i) detecting a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e)

in a sample from said subject;

and

(ii) diagnosing, or prognosing said neurological disease, wherein a varied level, or varied activity, or both said varied level and said varied activity, of at least one substance which is selected from the group consisting of a) to e) compared to a reference value representing a known health status;

or a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e) similar or equal to a reference value representing a known disease status indicates diagnosis, or prognosis of said neurological disease.

In preferred embodiments, said subjects suffer from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

It is particularly preferred that said sample is a brain tissue or other body cells including blood cells and skin fibroblasts. According to the present invention, a reduction in the level, or activity, or both said level and said activity, of *SELADIN-1* transcripts in said subject's brain regions affected heavily by neurodegeneration relative to a reference value indicates a diagnosis or prognosis of Alzheimer's disease. Predominantly neurons

within the inferior temporal lobe, the entorhinal cortex, the hippocampus and the amygdala degenerate in Alzheimer's disease.

It is also particularly preferred that said sample is cerebrospinal fluid or another body fluid.

It might be preferred that said subject has previously been determined to have one or more factors indicating that such subject is afflicted with a neurological disease.

In preferred embodiments, said subject can be a human, an experimental animal, e.g. a rat or a mouse, a domestic animal, or a non-human primate, e.g. a monkey. The experimental animal can be an animal model for a disorder, e.g. a transgenic mouse with an Alzheimer's-type neuropathology.

In preferred embodiments, at least one of said substances a) to e) is detected using an immunoassay, an enzyme activity assay and/or a binding assay.

In preferred embodiments, measurement of the level of transcription products of the *SELADIN-1* gene, or a variant thereof, is performed in body cells using Northern blots with probes specific for the *SELADIN-1* gene or said variant. Quantitative PCR with primer combinations to amplify *SELADIN-1* gene-specific sequences from cDNA obtained by reverse transcription of RNA extracted from body cells of a subject can also be applied. These techniques are known to those of ordinary skill in the art (see e.g. Watson et al., *Rekombinierte DNA*, 2nd edition, Spektrum Akademischer Verlag GmbH, Heidelberg, 1993; Watson et al., *Recombinant DNA*, 2nd ed. W.H. Freeman and Company, 1992).

In preferred embodiments, said level or activity of the protein molecule shown in SEQ ID NO. 1, or a variant or fragment thereof, is detected using an immunoassay. These assays can measure the amount of binding between said protein molecule ("SELADIN-1") and an anti-SELADIN-1 antibody by the use of enzymatic, chromodynamic, radioactive, or luminescent labels which are attached to either the anti-SELADIN-1

antibody or a secondary antibody which binds the anti-SELADIN-1 antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

The antibody or ligand to be used should preferably specifically detect SELADIN-1 or a variant or fragment thereof. It is preferred that it does not substantially interact with any other protein present in said sample.

Monoclonal antibodies capable of recognizing a protein molecule of SEQ ID NO. 1 or a variant or fragment thereof can be prepared using methods known in the art (see e.g. Köhler and Milstein, *Nature* 256, 495 - 497 1975; Kozbor et al., *Immunol. Today* 4, 72, 1983; Cole et al., *Monoclonal antibodies and cancer therapy*, Alan R. Liss, Inc., pp 77 - 96, 1985; Marks et al., *J. Biol. Chem.*, 16007 - 16010, 1992; the contents of which are incorporated herein by reference). Such monoclonal antibodies or fragments thereof can also be produced by alternative methods known to those of skill in the art of recombinant DNA technology (see e.g. Sastry et al, *PNAS* 86: 5728, 1989; ; Watson et al., *Rekombinierte DNA*, 2nd ed., Spektrum Akademischer Verlag GmbH, 1993; Watson et al, *Recombinant DNA*, 2nd ed., W. H. Freeman and Company, 1992; the contents of which are incorporated herein by reference). Monoclonal antibodies useful in the methods of the invention are directed to an epitope of SELADIN-1 or a variant or fragment thereof, such that the complex formed between the antibody and SELADIN-1, or between the antibody and said variant or fragment, can be recognized in detection assays. The term "antibodies" encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, single chain antibodies as well as fragments thereof which specifically bind to SELADIN-1, or to a variant or fragment thereof.

Antibodies or ligands might also be used in detecting specifically any molecules mentioned under d) to e) above.

If luminescent labels are used in any detection assay, it is preferred to use a confocal optical set-up.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to e) in a sample from a subject not suffering of said neurological disease. The healthy subject can be of the same weight, age, and gender as the subject who is being diagnosed or prognosed for said neurological disease. In some cases, it might be preferred to use a reference value from the subject which is diagnosed.

In another aspect, the kit may be used in monitoring a progression of said neurological disease in a subject.

In a further aspect, the kit may be used in monitoring success or failure of a therapeutic treatment of said subject.

In another aspect, the invention features a method of treating or preventing a neurological disease in a subject comprising administering to said subject in a therapeutically effective amount an agent or agents which directly or indirectly affect a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
- e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

In preferred embodiments, the method comprises the application of per se known methods of gene therapy and/or antisense nucleic acid technology to administer said agent or said agents.

In general, gene therapy includes several approaches: molecular replacement of a mutated gene, addition of a new gene resulting in the synthesis of a therapeutic protein, and modulation of endogeneous cellular gene expression by recombinant expression methods or by drugs. Gene-transfer techniques are described in detail (see e.g. Behr, Acc. Chem. Res. 26, 274 - 278, 1993; Mulligan, Science 260, 926 - 931, 1993; the contents of which are incorporated herein by reference) and include direct gene-transfer techniques such as mechanical microinjection of DNA into a cell as well as indirect techniques employing biological vectors (like recombinant viruses, especially retroviruses) or model liposomes, or techniques based on transfection with DNA coprecipitation with polycations, cell membrane perturbation by chemical (solvents, detergents, polymers, enzymes) or physical means (mechanic, osmotic, thermic, electric shocks). The postnatal gene transfer into the central nervous system has been described in detail (see e.g. Wolf, Current Opinion in Neurobiology, 3, 743 - 748, 1993; the contents of which are incorporated herein by reference).

In particular, the invention features a method of treating or preventing a neurological disease, such as Alzheimer's disease, by means of antisense nucleic acid therapy, i.e. the down-regulation of an inappropriately expressed or defective gene by the introduction of antisense nucleic acids or derivatives thereof into certain critical cells (see e.g. Gillespie, DN & P 5(7), 389 - 395, 1992; Agrawal, Tibtech 13, 197 - 199, 1996; Crooke, Bio/Technology 10, 882 - 886, 1992; the contents of which are incorporated herein by reference). Apart from hybridization strategies, the application of ribozymes, i.e. RNA molecules that act as enzymes, destroying RNA that carries the message of disease has also been described (see e.g. Barinaga, Science, 262, 1512 - 1514, 1993; the contents of which are incorporated herein by reference). It is particularly preferred that cells of the central nervous system, preferably the brain, of a subject are treated in such a way. Cell penetration can be performed by known strategies such as coupling of antisense nucleic acids and derivatives thereof to carrier particles, or the above

described techniques. Strategies for administering targeted therapeutic oligodeoxynucleotides are known to those of skill in the art (see e.g. Wickstrom, Tibtech, 10, 281 - 287, 1992; the contents of which are incorporated herein by reference). In some cases, delivery can be performed by mere topical application. Further approaches are directed to intracellular expression of antisense RNA. In this strategy, cells are transformed *ex vivo* with a recombinant gene that directs the synthesis of an RNA that is complementary to a region of the target nucleic acid. Therapeutically use of intracellularly expressed antisense RNA is procedurally similar to gene therapy.

In preferred embodiments, the method comprises grafting donor cells into the central nervous system, preferably the brain, of said subject, said subject or donor cells preferably treated so as to minimize or reduce graft rejection, wherein said donor cells are genetically modified by insertion of at least one transgene encoding said agent or agents. Said transgene might be carried by a viral vector, in particular a retroviral vector. The transgene can be inserted into the donor cells by a nonviral physical transfection of DNA encoding a transgene, in particular by microinjection. Insertion of the transgene can also be performed by electroporation, chemically mediated transfection, in particular calcium phosphate transfection, liposomal mediated transfection, etc.

In preferred embodiments, said agent is a therapeutic protein which can be administered to said subject, preferably a human, by a process comprising introducing subject cells into said subject, said subject cells having been treated *in vitro* to insert a DNA segment encoding said therapeutic protein, said subject cells expressing *in vivo* in said subject a therapeutically effective amount of said therapeutic protein. Said DNA segment can be inserted into said cells *in vitro* by a viral vector, in particular a retroviral vector.

In preferred embodiments, the therapeutic nucleic acid or protein reduces or prevents the degeneration of neurons and slows brain amyloid formation.

In another aspect, the invention features an agent which directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
- e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

In another aspect, the invention features a medicament comprising such an agent.

In still another aspect, the invention features an agent for treating or preventing a neurological disease, which agent directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
- e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

In preferred embodiments, said neurological diseases are Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell

degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

In a further aspect, the invention features the use of an agent, for treating or preventing a neurological disease, which agent directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
- e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

In preferred embodiments, said neurological diseases are Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death. Further examples of neurological diseases are Parkinson's disease, Huntington disease, Amyotrophic lateral sclerosis, Pick's disease.

In a further aspect, the invention features a method for identifying an agent that directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of

- a) a protein molecule shown in SEQ ID NO.1,
- b) a variant or fragment thereof,
- c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c).

e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c),

comprising the steps of:

- (i) providing a sample containing at least one substance which is selected from the group consisting of a) to e);
- (ii) contacting said sample with at least one agent;
- (iii) comparing an activity, or level, or both said activity and level, of at least one of said substances before and after said contacting.

Figure 1 depicts the selective vulnerability of brain regions in Alzheimer's disease. Predominantly neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus and the amygdala degenerate in Alzheimer's disease (Terry et al., *Annals of Neurology*, 10, 184-192, 1981). These brain regions are predominantly involved in the processing of learning and memory functions. In contrast, neurons within the frontal cortex, the occipital cortex and the cerebellum are largely intact and preserved from the neurodegenerative process in Alzheimer's disease.

Figure 2 discloses the identification of genes differentially expressed in brain regions from Alzheimer's disease patients. Brain areas with massive neuronal cell loss as well as areas with largely preserved neurons were identified and RNA extracted. Synthesis of cDNA was performed using an oligo-dT primer followed by PCR using the oligo-dT primer in combination with random primers and ($\alpha^{35}\text{S}$)-dATP. Reactions were separated on DNA sequencing gels, DNA bands visualized by autoradiography and bands lighting up in different intensities were cut out. DNA fragments were reamplified by PCR, cloned in *E. coli* and sequences determined. Expression and functional analyses were performed.

Figure 3 depicts the specifications of Alzheimer's disease brain tissue as it was used in the examples. Brain tissues from Alzheimer's disease patients and control subjects were removed within 6 hours of death, and immediately frozen on dry ice. For RNA extraction tissue sections from the inferior temporal lobe and frontal cortex were used.

Figure 4 discloses the quantification of *SELADIN-1* transcripts in brain tissue from Alzheimer's disease and control subjects by Northern blot analyses. Transcript levels were significantly lower in brain regions with severe neurodegeneration, i. e. temporal lobe in Alzheimer's disease (AD1-3) but not in normal brain (NB1-3), as compared to protected brain regions, i. e. frontal lobe. This decrease was specific as indicated by unchanged β -actin transcript levels used to control for equal loading of RNA.

Figure 5 depicts the transcription levels of the *SELADIN-1* gene in different human brain regions. The *SELADIN-1* gene was found to be expressed throughout the human brain. In particular transcription levels are high in all cortical areas, the hippocampus, the amygdala, the spinal cord, and the medulla. Note the unchanged levels in temporal lobe versus frontal lobe in this brain derived from a cognitively normal control subject without any signs of Alzheimer's disease. The analysis of β -actin transcripts was used as loading control.

Figure 6 depicts the distribution of *SELADIN-1* transcripts in human tissues. Comparable samples of RNA were spotted on nitrocellulose filters and *SELADIN-1* transcripts were quantified by hybridization using a labeled *SELADIN-1* gene specific probe. Significant levels of *SELADIN-1* gene transcripts were found in all brain regions tested. Transcripts were also detected in other tissues, however, strong variations in signal intensity indicated a tissue specific regulation of *SELADIN-1* expression.

Figure 7 depicts the expression of the *SELADIN-1* gene in rat brain cortex, hippocampus and basal nucleus analyzed by in situ hybridization. This staining pattern along with the higher magnifications indicate that *SELADIN-1* is predominantly expressed in neurons. No significant hybridization signals were observed with glial cells.

Figure 8 depicts the expression of *SELADIN-1* in rat brain nuclei. Strong *SELADIN-1* expression was found in the oculomotor, paraventricular, red and facial nuclei. Higher magnifications indicate predominant hybridization with neurons. No significant hybridization signals were observed with glial cells.

Figure 9 depicts the expression of *SELADIN-1* in rat brain hippocampus and substantia nigra. In situ hybridization with *SELADIN-1* transcripts was detected by photoemulsion-autoradiography, confirming the neuron specific expression of this gene.

Figure 10 discloses the subcellular localization of a *SELADIN-1*-EGFP (enhanced green fluorescent protein) fusion in transfected Cos cells. The confocal micrographs show the co-localization of the *SELADIN-1*-EGFP fusion with the golgi specific stain BODIPY TR ceramide indicating localization of *SELADIN-1* in the Golgi apparatus and the endoplasmic reticulum.

Figure 11 discloses that the *SELADIN-1*-EGFP fusion does not localize to mitochondria in transfected cos cells in spite of a putative mitochondrial targeting sequence close to the N-terminus of the *SELADIN-1* protein. The confocal micrographs show the different staining patterns caused by the *SELADIN-1*-EGFP fusion and the specific mitochondrial stain Mito Tracker Red CM-H₂XRos.

Figure 12 discloses structural features of the *SELADIN-1* protein based on multiple sequence alignments and secondary structure predictions. Near the N-terminus the *SELADIN-1* protein contains a putative mitochondrial localization signal that appears to be inactive in transfected Cos cells or when used in EGFP fusions. The central region of the protein contains a sequence that is homologous to a family of oxidoreductases and that contains a FAD site for covalent binding. The protein is predicted to contain five transmembrane regions. The expression in neurons, the co-localization in the Golgi apparatus and the endoplasmic reticulum of the *SELADIN-1* protein, the amyloid precursor protein (APP) and the presenilins PS1 and PS2 and furthermore the transmembrane character suggest a functional relationship between these proteins. Mutations in both APP and presenilins were shown to cause an increase in the production of β -amyloid. In a similar way the *SELADIN-1* protein might be involved in common biological pathways influencing the processing of the amyloid precursor protein and the generation of A β . Using the *SELADIN-1* protein as a probe, interaction partners can be identified which might represent new AD drug targets.

Figure 13 discloses the protein sequence of SELADIN-1 (SEQ ID NO. 1). The full length protein consists of 516 amino acid residues. The sequence is given in the one letter amino acid code.

Figure 14 discloses the nucleotide sequence of the cloned *SELADIN-1* cDNA (SEQ ID NO. 2) comprising 4248 nucleotides. The coding sequence for the SELADIN-1 protein starts at nucleotide position 100 and stops at position 1648.

Figure 15 discloses the comparison of nucleotide sequences of the cloned *SELADIN-1* cDNA comprising 4248 nucleotides and the *KIAA0018* cDNA comprising 4186 nucleotides. A significant difference exists at position 1228 of the *SELADIN-1* sequence where a C nucleotide (C/G basepair) is missing in the *KIAA0018* sequence. This results in a frameshift in the open reading frame in the *KIAA0018* sequence relative to the *SELADIN-1* sequence. The consequence is that the translation product of the *KIAA0018* gene is 390 amino acids in length compared to 516 amino acid residues of the *SELADIN-1* translation product. In addition to the difference in length, the frameshift causes a difference between the C-terminal 14 amino acids of the *KIAA0018* protein and the corresponding sequence area of the *SELADIN-1* polypeptide (pos. 377 - 390). The coding sequence for the SELADIN-1 protein starts at nucleotide position 100 and stops at position 1648.

EXAMPLE I

Post-mortem Alzheimer's disease brain tissues

Brain tissues from Alzheimer's disease patients and control subjects were removed within 6 hours of death, and immediately frozen on dry ice. Parallel sections were fixed in formaldehyde for histopathological confirmation of the diagnosis and for cell counts. Brain areas with massive neuronal cell loss as well as areas with largely preserved neurons were identified for comparisons of gene expression and stored at -80°C until RNA extractions were performed.

Identification of *SELADIN-1* by differential display PCR

Total RNA from post-mortem brain tissues was prepared by using the RNeasy kit (Qiagen). The RNA preparations were treated with DNase I (Boehringer Mannheim) together with RNasin (Promega) for 30 minutes, followed by phenol extraction, and ethanol precipitation. 0.2 mg of each RNA preparation were transcribed to cDNA by using Expand Reverse Transcriptase (Boehringer Mannheim) with one base anchor primers HT11A, HT11C and HT11G. In the following PCR reaction, the cDNAs were amplified by using HT11A along with the random primers HAP-5 (5'-TGCCGAAGCTTGGAGCTT-3') and HAP3-T (5'TGCCGAAGCTTTGGTCAT-3'). Taq-polymerase (AmpliTaq, Perkin Elmer Corp.), dGTP, dCTP, and dTTP (Amersham Pharmacia Biotech) and ($\alpha^{35}\text{S}$)-dATP (NEN life science products) were used in a PCR protocol according to Zhao et al. The PCR products were separated on 6% polyacrylamide-urea sequencing gels that were dried subsequently on 3 mm filter paper (Whatman), and X-ray films (Dupont) were exposed for 12 hours.

Cloning and sequencing

Differential bands were excised from the gel, boiled in water for 10 minutes, centrifuged, and cDNAs were precipitated from the supernatant fluids by using ethanol and glycogen/sodiumacetate, followed by dialysis against 10% glycerol for 1 hour through 0.025 mm filters (type VS, Millipore). The dialysates were used as templates for the reamplification reactions that were done under identical conditions as in the differential

display PCR, with the exception of the initial cycle for nonspecific annealing. The resulting PCR products were separated by agarose gelelectrophoresis, purified from the gel with the QIAEXII Agarose Gel Extraction Kit (Qiagen), and cloned into the *Hind* III restriction site of pBluescript KS (Stratagene). Cloned cDNA fragments were sequenced with an ABI 377 DNA sequencer (Perkin Elmer Corp.) by using T3 and T7 primers.

Amplification of a *SELADIN-1* cDNA-fragment

A *SELADIN-1* cDNA fragment was amplified by using cDNA transcribed from human brain tissue by using RNA High Fidelity Taq-polymerase (Boehringer Mannheim) and *SELADIN-1*-specific primers for a PCR reaction with 40 cycles of annealing at 70 °C for 1 minute, and polymerization at 72 °C for 3 minutes. The PCR products were separated by agarose gel electrophoresis, purified, and cloned into the *Sma* I restriction site of pBluescript KS (Stratagene). The cloned *SELADIN-1* PCR product was sequenced, and restriction fragments were used as probes both for screening a human brain cDNA library and for probing Northern blots.

Northern blotting

Total RNA from post-mortem human brains were prepared by using the Trizol reagent (Gibco BRL, Life Technologies), following the manufacturer's instructions. 5 - 10 mg of RNA were separated in 1 % formaldehyde-containing agarose gels, and the RNA was blotted onto nylon membranes (Hybond-N⁺, Amersham). Membranes were hybridized with (α -³²P)-dCTP (NEN) labeled *SELADIN-1*-specific cDNA probes that were generated by using the Megaprime DNA labelling kit (Amersham). Membranes were washed under high stringency conditions, and X-ray films were exposed for 1 to 72 hours. To control for equal loading of RNA, the identical membranes were probed with a 700 bp cDNA fragment of human glyceraldehyd-3-phosphate dehydrogenase (*GAPDH*), or with a β -actin cDNA fragment (Clontech).

In situ hybridization

Several *SELADIN-1*-specific cDNA probes of 650bp and of 900bp representing the initial two parts of the open reading frame were cloned in pBluescript (Stratagene) and reversely transcribed in the presence of (α^{35} S)-CTP by using the Ambion transcription kit. In situ hybridization was done with, 14mm sections of adult rat brain cut on a cryomicrotome, mounted on aminoalkylsilane-treated slides and fixed in 4 % paraformaldehyde in PBS for 5 min at room temperature. After washing for 5 min in PBS, sections were acetylated for 10 min, passed through a series of increasing ethanol grades and air dried. Prehybridizations were done in 50 % deionized formamide, 25 mM EDTA, 25 mM Pipes, pH 6.8, 0.75 M NaCl, 0.2 % SDS, 5xDenhardt's, 10 mM DTT, 250 mg/ml denatured herring sperm DNA and 250 mg/ml yeast tRNA. Hybridization of slides with RNA sense and antisense probes diluted to 2000 – 5000 cpm/ml in the same buffer with additional 10 % dextranulphate was performed at 50°C for 12 hours. Slides were then washed four times in 4xSSC for 5 min, each, followed by an incubation for 30 min. at 37 °C with 40 mg/ml RNaseA in 0.5 M NaCl, 10 mM Tris-HCL, pH 7.5, 1mM EDTA and another 30 min without RNaseA. Then slides were washed twice for 15 min. at 50°C in 2xSSC and dried through graded ethanols. Slides were exposed to Kodak Biomax x-ray films for 15 days and subsequently dipped in Kodak NTB-3 nuclear track emulsion and exposed for 6 weeks. After developing in Kodak D19 and fixing in Kodak Unifix, slides were counterstained with Mayer's Hemalaun and coverslipped.

Recombinant expression of SELADIN-1-EGFP fusion proteins in tissue culture

The complete coding region of *SELADIN-1* was subcloned into the N-terminus of the pEGFP-N1-expression vector (Clontech). Cos-7-cells were transfected with *EGFP* or with *SELADIN-1-EGFP* by using the SuperFect transfection reagent from Qiagen according to the manufacturers instructions. Cells were cultured in 3 cm dishes for two days. Part of the cells were stained for the Golgi-apparatus with 0.25 mM BODIPY TR ceramide (molecular probes) for one hour, the other part was treated with 250 nM of the mitochondrial stain Mito Tracker Red CM-H2Xros (Molecular probes) for 45 min. the subcellular localization of the *SELADIN-1-EGFP* fusion protein was analyzed by confocal laser scanning microscopy using the appropriate filter sets.

CLAIMS

1. An isolated nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1.
2. A DNA molecule having a coding sequence as shown in SEQ ID NO. 2 or a coding sequence substantially the same as the coding sequence of SEQ ID NO. 2.
3. A protein molecule shown in SEQ ID NO. 1.
4. A method of diagnosing or prognosing a neurological disease in a subject comprising:
determining a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of
 - a) a protein molecule shown in SEQ ID NO. 1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, a variant or fragment thereof,
 - d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
 - e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c)in a sample from said subject;
and
comparing said level, or said activity, or both said level and said activity, of at least one of said substances a) to e) to a reference value representing a known disease or health status,
thereby diagnosing or prognosing said neurological disease in said subject.

5. The method according to claim 4, wherein said subjects suffers from Alzheimer's disease and related neurofibrillary disorders, or neurodegenerative states characterized by cell degeneration or cell death.
6. A method of evaluating a treatment for a neurological disease, comprising:
determining a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of
 - a) a protein molecule shown in SEQ ID NO. 1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO. 1, a variant or fragment thereof,
 - d) a molecule affecting directly or indirectly a level, or an activity, or both said level and said activity, of at least one substance which is selected from the group consisting of a) to c),
 - e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c) in a sample from said subject;and
comparing said level, or said activity, or both said level and said activity, of at least one of said substances a) to e) to a reference value representing a known disease or health status,
thereby evaluating said treatment for said neurological disease.
7. An agent which directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of
 - a) a protein molecule shown in SEQ ID NO.1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,

- d) a molecule affecting directly or indirectly a level, or an activity, or both said level and activity, of at least one substance which is selected from the group consisting of a) to c).
 - e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).
8. A medicament comprising an agent according to claim 7.
9. An agent for treating or preventing a neurological disease, which agent directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of
- a) a protein molecule shown in SEQ ID NO.1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,
 - d) a molecule affecting directly or indirectly a level, or an activity, or both said level and activity of at least one substance which is selected from the group consisting of a) to c),
 - e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).
10. A medicament for treating or preventing a neurological disease comprising an agent according to claim 9.
11. Use of an agent for treating or preventing a neurological disease, which agent directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of
- a) a protein molecule shown in SEQ ID NO.1,
 - b) a variant or fragment thereof,
 - c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,

d) a molecule affecting directly or indirectly a level, or an activity, or both said level and activity of at least one substance which is selected from the group consisting of a) to c),

e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c).

12. A method for identifying an agent that directly or indirectly affects an activity, or level, or both said activity or level, of at least one substance which is selected from the group consisting of

a) a protein molecule shown in SEQ ID NO.1,

b) a variant or fragment thereof,

c) a nucleic acid molecule encoding a protein molecule shown in SEQ ID NO.1, a variant or fragment thereof,

d) a molecule affecting directly or indirectly a level, or an activity, or both said level and activity of at least one substance which is selected from the group consisting of a) to c),

e) a molecule which is affected in its level, or its activity, or both its level and activity, by at least one substance which is selected from the group consisting of a) to c)

comprising the steps of:

- (i) providing a sample containing at least one substance which is selected from the group consisting of a) to e);
- (ii) contacting said sample with at least one agent;
- (iii) comparing an activity, or level, or both said activity and level, of at least one of said substances before and after said contacting.

Selective vulnerability of brain regions in Alzheimer's disease





Figure 1

Identification of gene differentially expressed in AD brain regions

- 2/21 -




Figure 2

Identification of genes differentially expressed in AD brain regions

Material: AD brain tissue
post mortem time intervall <6h
2 different regions histologically characterized

- inferior temporal lobe
- frontal cortex

Method: mRNA differential display screen

Figur 3

Expression of Seladin-1 in AD brain

Additional RNA samples
from normal (NB) and AD brain

RNA samples used
for the DD screen




Figure 4

Expression of Seladin-1 in different human brain regions




Figure 5

Expression of Seladin-1 in human tissues

whole brain	amygdala	caudate nucleus	cerebellum	cerebral cortex					
occipital lobe	putamen			thalamus	sub-thalamic nucleus				
heart	aorta	skeletal muscle	colon	bladder	uterus				stomach
testis	ovary	pancreas	pituitary gland		thyroid gland				mammary gland
kidney		small intestine		thymus	peripheral leukocyte				bone marrow
appendix		trachea	placenta						
fetal brain	fetal heart	fetal kidney			fetal thymus				
young adult RNA 100ng	young adult RNA 100ng	E. coli RNA 100ng	E. coli RNA 100ng	Poly r(A) 100ng	human Ccst(DNA) 100ng	human DNA 100ng	human DNA 500ng		

Figure 6

Expression of Seladin-1 in rat brain




Figure 7

- 8/21 -


Expression of Seladin-1 in rat brain



Figure 8

- 9/21 -

Expression of Seladin-1 in rat brain



substantia nigra

hippocampus



Figure 9

Subcellular localization of Seladin-1 EGFP fusionprotein

Seladin-1-EGFP

Golgi stain
BODIPY TR ceramide

overlay



- 10/21 -

Figure 10

Subcellular localization of Seladin-1 EGFP fusionprotein

Seladin-1-EGFP

Mitochondrial stain
MitoTracker Red CM-H2XRos

overlay






Figure 11

- 11/21 -

- 12/21 -

Multiple sequence alignments and secondary structure prediction of Seladin-1




Figure 12

- 13/21 -
FIGURE 13: SEQ ID NO.1
Seladin-1 amino acid sequence

Seladin-1.orf Length: 516 May 29, 1998 14:51 Type: P Check: 1354 ..

1 MEPAVSLAVC ALLFLLWVRL KGLEFVLIHQ RWVPVCLFLL PLSLIFDIYY
51 YVRAWVVFKL SSAPRLHEQR VRDIQKQVRE WKEQGSKTFM CTGRPGWLTV
101 SLRVGKYKKT HKNIMINLMD ILEVDTKKQI VRVEPLVTMG QVTALLTSIG
151 WTLPVLPELD DLTVGGLIMG TGISSSSHKY GLFQHICTAY ELVLADGSFV
201 RCTPSENSDL FYAVPWSCGT LGFLVAAEIR IIPAKKYVKL RFEPVRGLEA
251 ICAKFTHESQ RQENHFVEGL LYSLDEAVIM TGVMTDEAEP SKLNSIGNYY
301 KPWFEEKHVEN YLKTNREGLE YIPLRHYYHR HTRSIFWELQ DIIPFGNNPI
351 FRYLFGWMVP PKISLLKLTO GETLRKLYEQ HHVVQDMLVP MKCLQQALHT
401 FQNDIHVYPI WLCPPFILPSQ PGLVHPKGNE AELYIDIGAY GEPRVKHFEA
451 RSCMRQLEKF VRSVHGFQML YADCYMNREE FWEMFDGSLY HKLREKLGCQ
501 DAFPEVYDKI CKAARH

- 14/21 -

FIGURE 14: SEQ ID NO.2

Seladin-1 cDNA sequence

Seladin-1 Length: 4248 April 28, 1998 14:10 Type: N Check: 8184 ..

```
1  cccgggctgt gggctacagg cgcagagcgg gccaggcgcg gagctggcgg
51  cagtgcacagg aggcgcgaac ccgcagcgcg taccgcgcgg cgcgcacca
101 tggagcccg cgtgtcgctg gccgtgtcg cgctgtctct cctgtgtgtg
151 gtgcgcctga aggggctgga gtccgtgtct atccaccagc gctgggtgtt
201 cgtgtgcctc ttcctcctgc cgctctcgct tatcttcgat atctactact
251 acgtgcgcgc ctgggtgggt ttcaagctca gcagcgcctc gcgcctgcac
301 gagcagcgcg tgcgggacat ccagaagcag gtgcgggaat ggaaggagca
351 gggtagcaag accttcattg gcacggggcg ccctggctgg ctactgtct
401 cactacgtgt cgggaagtac aagaagacac acaaaacat catgatcaac
451 ctgacggaca ttctggaagt ggacaccaag aaacagattg tccgtgtgga
501 gcccttgggt accatggggc aggtgactgc cctgtgacc tccattggct
551 ggactctccc cgtgttgctt gagcttgatg acctcacagt ggggggcttg
601 atcatgggca caggcatcga gtcatcatcc cacaagtacg gcctgttcca
651 acacatctgc actgcttacg agctggctct ggctgatggc agctttgtgc
701 gatgcactcc gtccgaaaac tcagacctgt tctatgccgt acctggctcc
751 tgtgggacgc tgggttctct ggtggccgct gagatccgca tcatccctgc
801 caagaagtac gtcaagctgc gtttcgagcc agtgcggggc ctggaggcta
851 tctgtgccaa gttcacccac ggtcccagc ggcaggagaa ccacttcgtg
901 gaagggtgc tctactccct ggatgaggct gtcattatga caggggtcat
951 gacagatgag gcagagccca gcaagctgaa tagcattggc aattactaca
1001 agcctgtggt ctttaagcat gtggagaact atctgaagac aaaccgagag
1051 ggcttgaggt acattccctt gagacactac taccaccgcc acacgcgcag
1101 catctctctg gagctccagg acatcatccc ctttggcaac aaccccatct
1151 tccgttacct ctttggctgg atggtgcctc ccaagatctc cctcctgaag
1201 ctgaccagg gtgagacctt gcgcaagctg tacgagcagc accacgtggt
1251 gcaggacatg ctggtgccca tgaagtgcct gcagcaggcc ctgcacacct
1301 tccaaaacga catccacgtc taccocatct ggctgtgtcc gttcatcctg
1351 cccagccagc caggcctagt gcaccccaaa ggaaatgagg cagagctcta
1401 catcgac tt ggagcatatg gggagccgcg tgtgaaacac ttgaaagcca
1451 ggtcctgcat gaggcagctg gagaagtttg tccgcagcgt gcatggcttc
1501 cagatgctgt atgccgactg ctacatgaac cgggaggagt tctgggagat
1551 gtttgatggc tccctgtacc acaagctgcg agagaagctg ggttgccagg
```

- 15/21 -

1601 acgccttccc cgaggtgtac gacaagatct gcaaggccgc caggcactga
1651 gctggagccc gcctggagag acagacacgt gtgagtggtc aggcattctc
1701 ccttcaactca agccttgcty ctttccctaga tccacacttt caaagagaaa
1751 cccctccaga actccacccc tgacagccca acaccacctt cctcctggct
1801 tccagggggc agcccagtg aatggaaaga atgtgggatt tggagtcaga
1851 caagcctgag tccagttccc cgtttagaac tcattagctg tgtgactctg
1901 ggtgagtcct ttaacccctc tgagcccggg tctcttcatt agttgaaagg
1951 gatagtaata cctacttgca gggtgtgtgc atctgagttg agcactggtc
2001 acattgaagg tgctgggtaa ggtgtagctc ttgttgcttc ccgttcagcg
2051 tcacatctgc agtggagcct gaaaaggctc cacattaggt caccgtgca
2101 cagccatggc tggaatgatg aaggggatac gctggagttg ccttgccatc
2151 gcctccatca gccagacgag gtccctcacag gagaaggaca gctcttcccc
2201 accctgggat ctccaggagg cagccacgga gtggggaggc ccagatgcy
2251 ctgtgccaaa gccaggctcg aggccaaagt tctccctgcc atccttggtg
2301 ccgtcctgcc ccttccctct tcatgctgg gcctgcaggc ccaccccagc
2351 caccactgag tccactcgga gtgcccctgt ttccctggaga aggcattcca
2401 ggggttgaatc ttgtcccagc ctccagcctgg gacacctagg tggagagagt
2451 ggtctccgct ctgaattgga tccaggggac ctgggctcat tcttcttggc
2501 tcaccaaccc tgcaggcctc atcttcccca aaacccactt tgtcttgggtg
2551 ggagtgggtc cgcgtgctc tgcagcaggg gctggggagt ggacagcatc
2601 aggtgggaaa gtggagtcca cccctcatgt tctgtaggat tctcaccgtg
2651 gggctgggaag aaaagagcat cgacttgatt tclccaacca ctcatccctc
2701 ttttcttttc tccaccact cccaccccca gctgtagtta atttcagtgc
2751 cttacaaatc ctaagctcag agaaagtccc atttccgttc cagagggag
2801 ggaacctccc taggtccttc cctggcttgt tataacgcaa agcttggttg
2851 tttatgcaac tctatcttaa gaactgccc gcctcagctg aaaacccgaa
2901 tctgagaagg aattgcgtca tgtaaggga gctggaatta agggagctga
2951 gccagtcatg gttgtggcgt gtgagtcagg agacctaggt ttcagccct
3001 ccttactgtc agcgagctgt gcaacgtggg caagtcattg tctctgagc
3051 tgcagtttcc tcatctgtca catcgctaca gacaagacct ccttggaaac
3101 cttctgattg tcttagacac tgtggttgca aaacccacgg aaagcctcat
3151 ttgtgtggaa agtcagagga aaaatgatcc agtggacact tggggattat
3201 ctgtcattca agatccttcc tccaacccca aggccagctc ccatctcatt
3251 tccagaaagg ctcatacctg gcttgcaggg aagcatctgt cttgtcattc
3301 caggtgccag aatcctctca gagtcatga aggtgttca cccatccac

- 16/21 -

3351 ccaaggettg gcacactgcc agtgtcttag cagggctctg tgagggctgg
3401 gggcatccag gcactcagaa ggcaaaggaa ccacctacc catttgacct
3451 ctggaggggg cagaagaaag aaagaaacct catcctatat ttacaaaagg
3501 atgtgaattc tggcattagc tctcatagga gacctatgtg cttccttgc
3551 cagtgcacaa ctgatgattc tacttgctgt agatgaatgg ttaacacgag
3601 ctagttaaac agtgccattg ttttgccagt gaagcctcca accctaagcc
3651 actgggacgg tggccagaga tgccagcagc ctctgtcgcc cttagtcata
3701 taacccaaat ccagacctta tccacaaccc ggggcttggg aaggaaggta
3751 ttttggaatc acacctccg gttatgttgc tccagtaaaa tcttgccctgg
3801 aaagaggcag tcttcttagc atggtgagct gagttcatgg ctttttttg
3851 tagccagtc tgtccctggc catccatgtg atggttttgg atggagttaa
3901 acttgatgcc agtgggcagt gcattgtggaa agtatcagag taagcctctc
3951 ccctccagag ccctgagttt cttggctgca tgaaggctct ctttagaatc
4001 agaattgtag ccagtttctt tggccagaag gatgaatact tggatattac
4051 tgaaggaggag gggcggagat ggggtgtggc gtgtatgggt tgtgattttt
4101 attttcttct ttggtcatgg gggccaagga gaaaggcatg aatcttccct
4151 gtcaggctct tacagccaca ggcactgtgt ctactgtctg gaagacatgt
4201 cccctggct gtggggccgc tgcttctgt taaataaag tggcctgg

- 17/21 -

FIGURE 15 CDNA sequence comparison
KIAA0018/Seladin-1

```
1  ggcggaaccgcgagcgcttaccgcgcggcgccgcaccatgg gcccgcc 50
62  ggcggaaccgcgagcgcttaccgcgcggcgccgcaccatggagcccgcc 111
51  gtgtcgctggccgtgtgcgcgctgctcttctgctgtgggtgcgcctgaa 100
112  gtgtcgctggccgtgtgcgcgctgctcttctgctgtgggtgcgcctgaa 161
101  ggggctggagttcggtgctcatccaccagcgctgggtgttcgtgtgectct 150
162  ggggctggagttcggtgctcatccaccagcgctgggtgttcgtgtgectct 211
151  tcctcctgcgctctcgcttatcttcgatatactactactacgtgcgcgcc 200
212  tcctcctgcgctctcgcttatcttcgatatactactactacgtgcgcgcc 261
201  tgggtggtgttcaagctcagcagcgctccgcgcctgcacgagcagcgct 250
262  tgggtggtgttcaagctcagcagcgctccgcgcctgcacgagcagcgct 311
251  gcgggacatccagaagcaggtgcgggaatggaaggagcagggtagcaaga 300
312  gcgggacatccagaagcaggtgcgggaatggaaggagcagggtagcaaga 361
301  ccttcagtgtgcacggggcgccctggctggctcactgtctcactacgtgtc 350
362  ccttcagtgtgcacggggcgccctggctggctcactgtctcactacgtgtc 411
351  ggggaagtacaagaagacacacaaaaacatcatgatcaacctgatggacat 400
412  ggggaagtacaagaagacacacaaaaacatcatgatcaacctgatggacat 461
401  tctggaagtggacaccaagaagacagattgtccgtgtggagcccttgggtga 450
462  tctggaagtggacaccaagaagacagattgtccgtgtggagcccttgggtga 511
451  ccatgggcccaggtgactgcccctgctgacctccattggctggactctcccc 500
512  ccatgggcccaggtgactgcccctgctgacctccattggctggactctcccc 561
501  gtgttgccctgagcttgatgacctcacagtggggggcttgatcatggggac 550
562  gtgttgccctgagcttgatgacctcacagtggggggcttgatcatggggac 611
551  aggcacgagtcacatcccacaagtaaggcctgttccaacacatctgca 600
612  aggcacgagtcacatcccacaagtaaggcctgttccaacacatctgca 661
601  ctgcttacgagctggtcctggctgatggcagctttgtgcatgcaactccg 650
662  ctgcttacgagctggtcctggctgatggcagctttgtgcatgcaactccg 711
651  tccgaaaactcagacctgttctatgccgtaccctggctcctgtgggacgct 700
712  tccgaaaactcagacctgttctatgccgtaccctggctcctgtgggacgct 761
701  ggggttctctgggtggccgctgagatccgcatcatccctgccaagaagtacg 750
762  ggggttctctgggtggccgctgagatccgcatcatccctgccaagaagtacg 811
751  tcaagctgcgtttcgagccagtgccggggcctggaggctatctgtgccaag 800
812  tcaagctgcgtttcgagccagtgccggggcctggaggctatctgtgccaag 861
801  ttcacccacgagtcacagcggcaggagaaccacttcgtggaagggtgct 850
862  ttcacccacgagtcacagcggcaggagaaccacttcgtggaagggtgct 911
```

- 18/21 -

851 ctactccctggatgaggctgtcattatgacaggggtcatgacagatgagg 900
|||||
912 ctactccctggatgaggctgtcattatgacaggggtcatgacagatgagg 961
|||||
901 cagagcccagcaagctgaatagcatttggcaattactacaagccgtgggttc 950
|||||
962 cagagcccagcaagctgaatagcatttggcaattactacaagccgtgggttc 1011
|||||
951 ttttaagcatgtggagaactatctgaagacaaaccgagagggcctggagta 1000
|||||
1012 ttttaagcatgtggagaactatctgaagacaaaccgagagggcctggagta 1061
|||||
1001 cattcccttgagacactactaccaccgccacacgcgcagcatcttcttggg 1050
|||||
1062 cattcccttgagacactactaccaccgccacacgcgcagcatcttcttggg 1111
|||||
1051 agctccaggacatcatcccccttggcaacaaccccatcttccgctacctc 1100
|||||
1112 agctccaggacatcatcccccttggcaacaaccccatcttccgctacctc 1161
|||||
1101 tttggctggatgggtgacctcccaagatctccctcctgaagctgacctcagg 1150
|||||
1162 tttggctggatgggtgacctcccaagatctccctcctgaagctgacctcagg 1211
|||||
1151 tgagaccctgcgcaag.tgtacgagcagcaccacgtgggtgcaggacatgc 1199
|||||
1212 tgagaccctgcgcaagctgtacgagcagcaccacgtgggtgcaggacatgc 1261
|||||
1200 tgggtgcccattgaagtgcctgcagcaggccctgcacaccttccaaaacgac 1249
|||||
1262 tgggtgcccattgaagtgcctgcagcaggccctgcacaccttccaaaacgac 1311
|||||
1250 atccacgtctacccccatctggctgtgtccgttcatectgccagccagcc 1299
|||||
1312 atccacgtctacccccatctggctgtgtccgttcatectgccagccagcc 1361
|||||
1300 aggcctagtgcacccccaaaggaaatgaggcagagctctacatcgacattg 1349
|||||
1362 aggcctagtgcacccccaaaggaaatgaggcagagctctacatcgacattg 1411
|||||
1350 gagcatatgggggagccgcgtgtgaaacactttgaagccaggctcctgcatg 1399
|||||
1412 gagcatatgggggagccgcgtgtgaaacactttgaagccaggctcctgcatg 1461
|||||
1400 aggcagctggagaagtttgtccgcagcgtgcatggcttccagatgctgta 1449
|||||
1462 aggcagctggagaagtttgtccgcagcgtgcatggcttccagatgctgta 1511
|||||
1450 tgccgactgctacatgaaccgggaggagttctgggagatgtttgatggct 1499
|||||
1512 tgccgactgctacatgaaccgggaggagttctgggagatgtttgatggct 1561
|||||
1500 ccttgtaccacaagctgagagagaagctgggttgccaggacgccttcccc 1549
|||||
1562 ccttgtaccacaagctgagagagaagctgggttgccaggacgccttcccc 1611
|||||
1550 gaggtgtacgacaagatctgcaaggccgcccaggcactgagctggagcccg 1599
|||||
1612 gaggtgtacgacaagatctgcaaggccgcccaggcactgagctggagcccg 1661
|||||
1600 cctggagagacagacacgtgtgagtggtcaggcatcttcccttcactcaa 1649
|||||
1662 cctggagagacagacacgtgtgagtggtcaggcatcttcccttcactcaa 1711
|||||
1650 gcttggctgcttttccatagatccacactttcaaagagaaaacccctccagaa 1699
|||||
1712 gcttggctgcttttccatagatccacactttcaaagagaaaacccctccagaa 1761
|||||

12-11-1998

- 19/21 -

1700 ctcccacccctgacagcccaacaccaccttccctcctggcttccagggggca 1749
1762 ctcccacccctgacagcccaacaccaccttccctcctggcttccagggggca 1811
1750 gcccagtggaatggaaagaatgtgggatttggagtcagacaagcctgagt 1799
1812 gcccagtggaatggaaagaatgtgggatttggagtcagacaagcctgagt 1861
1800 ccagttccccggttagaactcattagctgtgtgactctgggtgagtcctt 1849
1862 ccagttccccggttagaactcattagctgtgtgactctgggtgagtcctt 1911
1850 taacccctctgagccccgggtctcttcattagttgaaagggatagtaatac 1899
1912 taacccctctgagccccgggtctcttcattagttgaaagggatagtaatac 1961
1900 ctacttgacaggttgttgtcatctgagttgagcactgggtcacattgaagg 1949
1962 ctacttgacaggttgttgtcatctgagttgagcactgggtcacattgaagg 2011
1950 gctgggtaagtggttagctcttgttgcctcccggttcagcgtcacatctgca 1999
2012 gctgggtaagtggttagctcttgttgcctcccggttcagcgtcacatctgca 2061
2000 gtggagcctgaaaaggctccacattaggtcacctgtgcacagccatgggt 2049
2062 gtggagcctgaaaaggctccacattaggtcacctgtgcacagccatgggt 2111
2050 ggaatgatgaaggggatacgttgagttgcccctgccatcgccctccatcag 2099
2112 ggaatgatgaaggggatacgttgagttgcccctgccatcgccctccatcag 2161
2100 ccagacgaggtcctcacaggagaaggacagctcttccccaccctggggtc 2149
2162 ccagacgaggtcctcacaggagaaggacagctcttccccaccctggggtc 2211
2150 tcaggaggggcagccacggagtggggagggcccgatgcgctgtgccaaag 2199
2212 tcaggaggggcagccacggagtggggagggcccgatgcgctgtgccaaag 2261
2200 ccaggtccgagggccaaagtcttccctgccatccttgggtgccgtcctgcc 2249
2262 ccaggtccgagggccaaagtcttccctgccatccttgggtgccgtcctgcc 2311
2250 cttctctcttcatgcttgggcctgcagggccacccagccaccactgagt 2299
2312 cttctctcttcatgcttgggcctgcagggccacccagccaccactgagt 2361
2300 ccactcggagtgccctgtgttccctggagaaggcattccaggggttgaatct 2349
2362 ccactcggagtgccctgtgttccctggagaaggcattccaggggttgaatct 2411
2350 tgtcccagcctcagcctgggacacctaggtggagagagtggtctccgctc 2399
2412 tgtcccagcctcagcctgggacacctaggtggagagagtggtctccgctc 2461
2400 tgaattggatccaggggacctggggtcattcttcttgggtcaccacccct 2449
2462 tgaattggatccaggggacctggggtcattcttcttgggtcaccacccct 2511
2450 gcaggcctcatcttccccaaaaccaccttcttcttgggtgggagtggggtcc 2499
2512 gcaggcctcatcttccccaaaaccaccttcttcttgggtgggagtggggtcc 2561
2500 gcgctgctctgcagcaggggctggggagtgagcagcatcaggtgggaaag 2549
2562 gcgctgctctgcagcaggggctggggagtgagcagcatcaggtgggaaag 2611

12-11-1998

EP98121478.6

SPEC

- 20/21 -

2550 tggagtcacacccatggtttctgtaggattctcaccgtggggctggaaga 2599
2612 tggagtcacacccatggtttctgtaggattctcaccgtggggctggaaga 2661
2600 aaagagcatcgacttgatttctccaaccactcatccctcttttctttct 2649
2662 aaagagcatcgacttgatttctccaaccactcatccctcttttctttct 2711
2650 tccaccactccccacccagctgtagttaatttcagtgccttacaaatcc 2699
2712 tccaccactccccacccagctgtagttaatttcagtgccttacaaatcc 2761
2700 taagctcagagaaagtccatttccgttccagaggggaagggaacctccct 2749
2762 taagctcagagaaagtccatttccgttccagaggggaagggaacctccct 2811
2750 aggtccctccctggcttggtataacgcaaagcttggttggttatgcaact 2799
2812 aggtccctccctggcttggtataacgcaaagcttggttggttatgcaact 2861
2800 ctatcttaagaactgccagcctcagctgaaaaccggaatctgagaagga 2849
2862 ctatcttaagaactgccagcctcagctgaaaaccggaatctgagaagga 2911
2850 attgcgtcatgtaagggaagctggaattaagggaagctgagccagtcattg 2899
2912 attgcgtcatgtaagggaagctggaattaagggaagctgagccagtcattg 2961
2900 ttgtggcggtgtgagtcaggagacctaggtttcagccctctctactgtca 2949
2962 ttgtggcggtgtgagtcaggagacctaggtttcagccctctctactgtca 3011
2950 gcgagctgtgcaacgtgggcaagctcattgtctctgagctgcagtttct 2999
3012 gcgagctgtgcaacgtgggcaagctcattgtctctgagctgcagtttct 3061
3000 catctgtcacatcgctacagacaagacctccctggaaccttctgattgt 3049
3062 catctgtcacatcgctacagacaagacctccctggaaccttctgattgt 3111
3050 cttagacactgtggttgcaaaacccacggaaagcctcatttgtgtgga 3099
3112 cttagacactgtggttgcaaaacccacggaaagcctcatttgtgtgga 3161
3100 gtcagagggaataatgatccagtggaacttggggattatctgtcattcaa 3149
3162 gtcagagggaataatgatccagtggaacttggggattatctgtcattcaa 3211
3150 gatccttcccttcaaccccaaggccagctcccatctcatttccagaaaggc 3199
3212 gatccttcccttcaaccccaaggccagctcccatctcatttccagaaaggc 3261
3200 tcatacctggcttgcaagggaagcatctgtcttgatccaggtgccaga 3249
3262 tcatacctggcttgcaagggaagcatctgtcttgatccaggtgccaga 3311
3250 atcctctcagagtcattgaagggtgttcacccatccacccaaggcttgg 3299
3312 atcctctcagagtcattgaagggtgttcacccatccacccaaggcttgg 3361
3300 cacactgccagtgctcttagcagggtcttgtgagggtgggggcatccagg 3349
3362 cacactgccagtgctcttagcagggtcttgtgagggtgggggcatccagg 3411
3350 cactcagaaggcaaaaggaaccacccctaccatttggcctctggagggggc 3399
3412 cactcagaaggcaaaaggaaccacccctaccatttggcctctggagggggc 3461

EP98121478.6

12-11-1998

- 21/21 -

3400 agaagaaagaaagaaacctcatcctatatattttacaaagcatgtgaattct 3449
|||||
3462 ag agaaagaaagaaacctcatcctatatattttacaaagcatgtgaattct 3511
|||||
3450 ggcattagctctcataggagacccatgtgcttccttgctcagtgcaaaac 3499
|||||
3512 ggcattagctctcataggagacccatgtgcttccttgctcagtgcaaaac 3561
|||||
3500 tgatgattctacttgctgtagatgaatggtaaacacgagctagttaaaca 3549
|||||
3562 tgatgattctacttgctgtagatgaatggtaaacacgagctagttaaaca 3611
|||||
3550 gtgccattgttttgccagtgaagcctccaaccctaagccactgggacggg 3599
|||||
3612 gtgccattgttttgccagtgaagcctccaaccctaagccactgggacggg 3661
|||||
3600 ggccagagatgccagcagcctctgtcgcccttagtcatataaccnaaac 3649
|||||
3662 ggccagagatgccagcagcctctgtcgcccttagtcatataaccnaaac 3711
|||||
3650 cagaccttatccacaaccggggcttggaaaggaaggatattttggaatca 3699
|||||
3712 cagaccttatccacaaccggggcttggaaaggaaggatattttggaatca 3761
|||||
3700 caccctccggttatgttgctccagtaaaatcttgccctggaaagaggcagt 3749
|||||
3762 caccctccggttatgttgctccagtaaaatcttgccctggaaagaggcagt 3811
|||||
3750 cttcttagcatgggtgagctgagttcatggctttttttgtagccagtcct 3799
|||||
3812 cttcttagcatgggtgagctgagttcatggctttttttgtagccagtcct 3861
|||||
3800 gtccctggccatccatgtgatgggttttgatggagttaaacttgatgcca 3849
|||||
3862 gtccctggccatccatgtgatgggttttgatggagttaaacttgatgcca 3911
|||||
3850 gtgggcagtgcatgtggaaagtatcagagtaagcctctccctccagagc 3899
|||||
3912 gtgggcagtgcatgtggaaagtatcagagtaagcctctccctccagagc 3961
|||||
3900 cctgagtttcttggtgcatgaagggttttctttagaatcagaattgtagc 3949
|||||
3962 cctgagtttcttggtgcatgaagggttttctttagaatcagaattgtagc 4011
|||||
3950 cagtttctttggccagaaggatgaataacttgatattactgaaagggagg 3999
|||||
4012 cagtttctttggccagaaggatgaataacttgatattactgaaagggagg 4061
|||||
4000 ggtggagatgggtgtggcagtgatgggtgtgatttttatttcttctt 4049
|||||
4062 ggtggagatgggtgtggcagtgatgggtgtgatttttatttcttctt 4111
|||||
4050 tggatcatgggggccaaggagaaaggcatgaatcttccctgtcaggctctt 4099
|||||
4112 tggatcatgggggccaaggagaaaggcatgaatcttccctgtcaggctctt 4161
|||||
4100 acagccacaggcactgtgtctactgtctggaagacatgtcccggtggctg 4149
|||||
4162 acagccacaggcactgtgtctactgtctggaagacatgtcccggtggctg 4211
|||||
4150 tggggccgctgcttctgttttaataaaaagtggcctgg 4186
|||||
4212 tggggccgctgcttctgttttaataaaaagtggcctgg 4248

Abstract

An isolated nucleic acid molecule encoding a protein molecule shown in SEQ ID
No. 1.

THIS PAGE BLANK (USPTO)